

In the Specification

At page 7, paragraph 16, please insert the following replacement paragraph.

[0016] The topical composition of the invention can contain additional ingredients commonly found in skin care compositions and cosmetics, such as, for example, tinting agents, emollients, skin conditioning agents, emulsifying agents, humectants, preservatives, antioxidants, perfumes, chelating agents, etc., provided that they are physically and chemically compatible with other components of the composition. Preservatives include, but are not limited to, C1-C3 alkyl parabens and phenoxyethanol, typically present in an amount ranging from about 0.5% to about 2.0% by weight percent, based on the total composition. Emollients, typically present in amounts ranging from about 0.01% to 5% of the total composition include, but are not limited to, fatty esters, fatty alcohols, mineral oils, polyether siloxane copolymers, and mixtures thereof. Humectants, typically present in amounts ranging from about 0.1% to about 5% by weight of the total composition include, but are not limited to, polyhydric alcohols such as glycerol, polyalkylene glycols (e.g., butylene glycol, propylene glycol, dipropylene glycol, polypropylene glycol, and polyethylene glycol) and derivatives thereof, alkylene polyols and their derivatives, sorbitol, hydroxy sorbitol, hexylene glycol, 1,3-dibutylene glycol, 1,2,6-hexanetriol, ethoxylated glycerol, propoxylated glycerol, and mixtures thereof. Emulsifiers, typically present in amounts from about 1% to about 10% by weight of the composition, include, but are not limited to, stearic acid, cetyl alcohol, stearyl alcohol, steareth 2, steareth 20, acrylates/C10-30 alkyl acrylate crosspolymers, and mixtures thereof. Chelating agents, typically present in amounts ranging from about 0.01% to about 2% by weight, include, but are not limited to, ethylenediamine tetraacetic acid (EDTA) and derivatives and salts thereof, dihydroxyethyl glycine, tartaric acid, and mixtures thereof. Antioxidants, typically present in an amount ranging from about 0.02% to about 0.5% by weight of the composition, include, but are not limited to, butylated hydroxy toluene (BHT); vitamin C and/or vitamin C derivatives, such as fatty acid esters of ascorbic acid, particularly ascorbyl palmitate; butylated hydroanisole (BHA); phenyl- α -

naphthylamine; hydroquinone; propyl gallate; nordihydroquiaretic acid; vitamin E and/or derivatives of vitamin E, including tocotrienol and/or tocotrienol derivatives; calcium pantothenates; green tea extracts; mixed polyphenols; and mixtures of any of these. As mentioned above, particularly preferred antioxidants are those that provide additional benefits to the skin such as ascorbyl palmitate. (See additional ingredients and methods in U.S. Pat. Nos. 4,775,530, 5,376,361, 5,409,693, 5,545,398, 5,574,063, 5,643,586, 5,709,868, 5,879,690, 5,965,618, ~~5,968,618~~, 6,051,244, 6,162,419, and 6,191,121 to Perricone).